

USN

--	--	--	--	--	--	--	--	--	--

17MAT41

## Fourth Semester B.E. Degree Examination, Dec.2019/Jan.2020 Engineering Mathematics - IV

Time: 3 hrs.

Max. Marks: 100

*Note: Answer any FIVE full questions, choosing ONE full question from each module.*

### Module-1

- 1 a. From Taylor's series method, find  $y(0.1)$ , considering upto fourth degree term if  $y(x)$  satisfying the equation  $\frac{dy}{dx} = x - y^2$ ,  $y(0) = 1$ . (06 Marks)
- b. Using Runge-Kutta method of fourth order  $\frac{dy}{dx} + y = 2x$  at  $x = 1.1$  given that  $y = 3$  at  $x = 1$  initially. (07 Marks)
- c. If  $\frac{dy}{dx} = 2e^x - y$ ,  $y(0) = 2$ ,  $y(0.1) = 2.010$ ,  $y(0.2) = 2.040$  and  $y(0.3) = 2.090$ , find  $y(0.4)$  correct upto four decimal places by using Milne's predictor-corrector formula. (07 Marks)

### OR

- 2 a. Using modified Euler's method find  $y$  at  $x = 0.2$  given  $\frac{dy}{dx} = 3x + \frac{1}{2}y$  with  $y(0) = 1$  taking  $h = 0.1$ . (06 Marks)
- b. Given  $\frac{dy}{dx} + y + zy^2 = 0$  and  $y(0) = 1$ ,  $y(0.1) = 0.9008$ ,  $y(0.2) = 0.8066$ ,  $y(0.3) = 0.722$ . Evaluate  $y(0.4)$  by Adams-Bashforth method. (07 Marks)
- c. Using Runge-Kutta method of fourth order, find  $y(0.2)$  for the equation  $\frac{dy}{dx} = \frac{y-x}{y+x}$ ,  $y(0) = 1$  taking  $h = 0.2$ . (07 Marks)

### Module-2

- 3 a. Apply Milne's method to compute  $y(0.8)$  given that  $\frac{d^2y}{dx^2} = 1 - 2y \frac{dy}{dx}$  and the following table of initial values.

x	0	0.2	0.4	0.6
y	0	0.02	0.0795	0.1762
y'	0	0.1996	0.3937	0.5689

- b. Express  $f(x) = x^4 + 3x^3 - x^2 + 5x - 2$  in terms of Legendre polynomials. (07 Marks)
- c. Obtain the series solution of Bessel's differential equation  $x^2y'' + xy' + (x^2 + n^2)y = 0$  leading to  $J_n(x)$ . (07 Marks)

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.  
2. Any revealing of identification, appeal to evaluator and /or equations written eg, 42+8 = 50, will be treated as malpractice.



OR

- 4 a. Given  $y'' - xy' - y = 0$  with the initial conditions  $y(0) = 1, y'(0) = 0$ , compute  $y(0.2)$  and  $y'(0.2)$  using fourth order Runge-Kutta method. (06 Marks)
- b. Prove  $J_{-1/2}(k) = \sqrt{\frac{2}{\pi x}} \cos x$ . (07 Marks)
- c. Prove the Rodrigues formula  $P_n(x) = \frac{1}{2^n n!} \frac{d^n y}{dx^n} (x^2 - 1)^n$  (07 Marks)

**Module-3**

- 5 a. Derive Cauchy-Riemann equations in Cartesian form. (06 Marks)
- b. Discuss the transformation  $w = z^2$ . (07 Marks)
- c. By using Cauchy's residue theorem, evaluate  $\int_C \frac{e^{2z}}{(z+1)(z+2)} dz$  if C is the circle  $|z| = 3$ . (07 Marks)

OR

- 6 a. Prove that  $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right) |f(z)|^2 = 4|f'(z)|^2$  (06 Marks)
- b. State and prove Cauchy's integral formula. (07 Marks)
- c. Find the bilinear transformation which maps  $z = \infty, i, 0$  into  $w = -1, -i, 1$ . (07 Marks)

**Module-4**

- 7 a. Find the mean and standard of Poisson distribution. (06 Marks)
- b. In an examination 7% of students score less than 35 marks and 89% of the students score less than 60 marks. Find the mean and standard deviation if the marks are normally distributed given  $A(1.2263) = 0.39$  and  $A(1.4757) = 0.43$  (07 Marks)
- c. The joint probability distribution table for two random variables X and Y is as follows:

	Y			
X	-2	-1	4	5
1	0.1	0.2	0	0.3
2	0.2	0.1	0.1	0

Determine:

- i) Marginal distribution of X and Y
- ii) Covariance of X and Y
- iii) Correlation of X and Y (07 Marks)


OR

- 8 a. A random variable X has the following probability function:

x	0	1	2	3	4	5	6	7
P(x)	0	K	2k	2k	3k	K <sup>2</sup>	2k <sup>2</sup>	7k <sup>2</sup> +k

Find K and evaluate  $P(x \geq 6), P(3 < x \leq 6)$ . (06 Marks)

- b. The probability that a pen manufactured by a factory be defective is 1/10. If 12 such pens are manufactured, what is the probability that
  - i) Exactly 2 are defective
  - ii) Atleast two are defective
  - iii) None of them are defective. (07 Marks)
- c. The length of telephone conversation in a booth has been exponential distribution and found on an average to be 5 minutes. Find the probability that a random call made
  - i) Ends in less than 5 minutes
  - ii) Between 5 and 10 minutes. (07 Marks)

**Module-5**

- 9 a. A die is thrown 9000 times and a throw of 3 or 4 was observed 3240 times. Show that the die cannot be regarded as an unbiased die. (06 Marks)
- b. A group of 10 boys fed on diet A and another group of 8 boys fed on a different diet B for a period of 6 months recorded the following increase in weight (lbs):

Diet A:	5	6	8	1	12	4	3	9	6	10
Diet B:	2	3	6	8	10	1	2	8		

Test whether diets A and B differ significantly  $t_{.05} = 2.12$  at 16df. (07 Marks)

- c. Find the unique fixed probability vector for the regular stochastic matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 1/6 & 1/2 & 1/3 \\ 0 & 2/3 & 1/3 \end{bmatrix}$$

(07 Marks)

OR

- 10 a. Define the terms:
- Null hypothesis
  - Type-I and Type-II error
  - Confidence limits

(06 Marks)

- b. The t.p.m. of a Markov chain is given by  $P = \begin{bmatrix} 1/2 & 0 & 1/2 \\ 1 & 0 & 0 \\ 1/4 & 1/2 & 1/4 \end{bmatrix}$ . Find the fixed probabilities vector. (07 Marks)

- c. Two boys  $B_1$  and  $B_2$  and two girls  $G_1$  and  $G_2$  are throwing ball from one to another. Each boy throws the ball to the other boy with probability  $1/2$  and to each girl with probability  $1/4$ . On the other hand each girl throws the ball to each boy with probability  $1/2$  and never to the other girl. In the long run how often does each receive the ball? (07 Marks)

\*\*\*\*\*