

Any revealing of identification, appeal to evaluator and /or equations written eg. 42+8=50, will be treated as malpractice. Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.



OR

- Prove that angle of friction is equal to angle to repose. a.
- Define: b.

b.

- i) Limiting friction
- ii) Cone of friction.
- Two blocks A and B of masses 30kg and 40kg respectively are connected by means of a bar c. placed on a inclined plane as shown in Fig.Q.4(c). If M = 0.25, find the value of  $\theta$  required for the motion of block to impend. (10 Marks)



# Module-3

- Explain the different types of loading with sketches. 5 a. Briefly explain determinate and indeterminate beams.
- (06 Marks) (04 Marks)
- A beam AB 1.7m long is loaded as shown in Fig.Q.5(c). Find the reactions at A and B. c.



- What are the steps to be followed in the analysis of statically determinate trusses by the 6 a. method of joints? (06 Marks)
  - Analyze the truss shown in Fig.Q.6(b) by the method of joints. Tabulate the results and b. indicate the nature of forces in the truss. (14 Marks)



## (04 Marks)

(06 Marks)

### 18CIV14/24

### **Module-4**

- State and prove parallel axis theorem. 7 a. (08 Marks) Find the centroid for the Fig.Q.7(b) shown below: b. ÆУ CENTRA IBRAR ₽ž 80mm 80mm + Fig.Q.7(b)(12 Marks) OR 8
  - Find the centroid of a triangle of base 'b' and height 'h' from first principle. (08 Marks) a. b.
    - Determinate the moment of inertia of Fig.Q.8(b) about centroidal X axis and Y axis.



- 9 Define: a.
  - i) Trajectory
    - Time of flight ii)
    - iii) Superelevation
    - iv) Curve linear motion
    - Rectilinear motion. v)
  - A tower in 90m in height. A particle is dropped from the top of tower and at the same time b. another particle is projected upward from the foot of tower. Both the particle meet at a height of 30m with respect foot of the tower. Find the velocity with which second particle is projected upward. (10 Marks)

### OR

- 10 State and explain D'Alembert's principle. a.
  - Two weights 800N and 200N are connected by thread and they move along a rough b. 🗖 horizontal plane under the action of force of 400N applied to the 800N weight as shown in Fig.Q.10(b). Using D'Alembert's principle, find the acceleration of weights and tension in thread. Take coefficient of friction = 0.3. (12 Marks)



#### (10 Marks)

(12 Marks)

### (08 Marks)